

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2007		2. REPORT TYPE		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Mass Casualties in Combat: Lessons Learned				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Medical Corps, Madigan Army Medical Center, Tacoma, WA, 98431				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Mass Casualties in Combat: Lessons Learned

MAJ Alec C. Beekley, MD, FACS

J Trauma. 2007;62:S39–S40.

The generation of *multiple* casualty events, where the number of casualties do not overwhelm medical resources, is the norm in modern warfare, and the current military trauma systems in Iraq and Afghanistan routinely and effectively deal with these events. *Mass* casualty events, where the number of casualties overwhelms the local medical resources, are less common events but still occur with greater frequency during military conflicts than in civilian settings. Recent terrorist attacks in civilian settings, such as the attacks on the World Trade Center on September 11, 2001, and the Madrid and London train bombings, have blurred the traditional lines between civilian and military trauma by generating large number of casualties at once with injuries from explosives. Hence, the experiences and lessons learned by military surgeons in modern conflicts with regard to mass casualty events undoubtedly resonate strongly with our civilian colleagues and must continue to be shared.

The military trauma system in Iraq has evolved from one based largely on small, mobile, forward surgical (echelon IIb) units during the maneuver phase of the war to a robust, well-planned system based primarily on larger fixed facilities (e.g. combat support hospitals, echelon III).¹ Forward surgical units currently augment the echelon III facilities or provide task-oriented surgical support in areas of high operation tempo. This allows for the evacuation of the majority of casualties directly from point of wounding to a facility where maximal surgical, medical, and blood bank support can be leveraged and damage control surgical techniques combined with damage control resuscitation can be brought to bear.²

Critical to the successful management of multiple and mass casualties is the process of triage. The Emergency War Surgery Manual provides important guidelines for this process, emphasizing that triage must occur at multiple levels of care and be repeated until all casualties are treated.³ Experiences in Iraq provide several lessons learned regarding triage, as follows: 1. proximity of hospital to mass casualty event

can complicate triage; 2. performance of triage is effected by casualties' modes of arrival, the physical layout of the medical treatment facility, and the presence of incoming hostile fire (among other factors); 3. triage and re-triage should occur at or near the point of wounding, on the helipad at receiving facilities, at the trauma bay door, in the trauma bays, and in the operating room; 4. radial pulse character and Glasgow Coma Scale Motor score can provide fast and reliable triage criteria for distinguishing between stable and unstable patients⁴; 5. initial triage at echelon III facilities should be performed by senior/experienced surgical personnel who are not directly involved in patient care; 6. providers assigned to specific patients should stay with those patients unless reassigned by triage officer; 7. Focused Abdominal Sonography for Trauma (FAST) and CT scan (where available) can be effective triage tools to prioritize patients^{5–7}; and 8. surgeons must be willing to re-triage patients in the operating room as expectant based on injuries identified, blood products required, physiologic status of patient, and number of other casualties requiring operation.

The ability to predict resource needs based on the number of casualties generated by a mass casualty event was described by Soffer et al. in their analysis of blood product utilization after terrorist attacks in Israeli.⁸ These researchers created a novel predictive index, the packed cells per patient index (PPI), and found that the average PPI for 19 consecutive terrorist attacks was 1.0. 5.7% of their casualties required massive transfusion (>10 units of blood). A similar analysis was applied to the management of 50 multiple or mass casualty events by a single combat support hospital (CSH) in Iraq. The number of casualties treated during these events was (at least) greater than one SD over the mean number of casualties treated per day at the CSH. The average number of patients for these 50 days was 24. 1327 units of blood were transfused on these days for an average of 26.5 units per day. On average, 20% of casualties required transfusion and 5% required massive transfusion. These percentages remained consistent regardless of number of casualties per event or predominance of blast or gunshot wounds as wounding mechanism, with the exception that when the mass casualty event was caused by a single explosion, the percentage of patients requiring massive transfusion doubled (9.6%). The average PPI for these days was 1.12. These numbers are strikingly similar to Soffer's findings. Days where the PPI greatly exceeded the average were rare, and analysis of these events revealed cases where casualties were over- or under-triaged with resultant expenditure of large amounts of blood products

Copyright © 2007 by Lippincott Williams & Wilkins, Inc.

This article was written for the proceedings from a conference entitled *12th Annual San Antonio Trauma Symposium* in San Antonio, Texas. The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

US Army Medical Corps, Madigan Army Medical Center, Tacoma, WA; email: alec.beekley@amedd.army.mil.

DOI: 10.1097/TA.0b013e31806541e9

on casualties who were likely unsalvageable. Finally, there was no correlation between the number of casualties per event and mortality.

In summary, the current military trauma systems in Iraq and Afghanistan are effective at dealing with both routine multiple casualty events and the less common mass casualty event. Analysis of multiple and mass casualty events from current conflicts can provide critical lessons learned regarding triage and resource utilization which can potentially be applied to other conflicts or civilian multiple or mass casualty events. Although the findings of this study regarding blood product utilization may not be directly applicable to catastrophic, national-level events, the consistency of the findings with the Israeli experience and across a wide range of smaller multiple or mass casualty explosion-related events is striking. Hence, for such events, the number of casualties generated may provide a reliable *baseline* prediction of blood product needs and percentages of patients who will require transfusion and massive transfusion. This in turn can help determine distribution of casualties among medical treatment facilities and allow for activation of additional blood resources, such as notification of more distant blood banks or initiation of fresh whole blood drives. Clearly, the experiences with mass ca-

sualty events that both military and civilian surgeons encounter must continue to be studied and shared.

REFERENCES

1. Beekley AC. United States military surgical response to modern large-scale conflicts: the ongoing evolution of a trauma system. *Surg Clin North Am.* 2006;86:689–709.
2. Holcomb JB. Damage Control Resuscitation: Directly Addressing the Coagulopathy of Trauma. Submitted, 2006.
3. Emergency War Surgery, 3rd United States Revision. 2004.
4. Holcomb JB, Salinas J, McManus JM, et al. Manual vital signs reliably predict need for life-saving interventions in trauma patients. *J Trauma.* 2005;59:821–828.
5. Udobi KF, Rodriguez A, Chiu WC, et al. Role of ultrasonography in penetrating abdominal trauma: a prospective clinical study. *J Trauma.* 2001;50:475–479.
6. Blaivas M. Triage in the trauma bay with the focused abdominal sonography for trauma (FAST) examination. *J Emerg Med.* 2001; 21:41–44.
7. Beekley A, Blackburne L, Sebesta J, et al. Evaluation for Intra-abdominal Injury from Combat Fragmentation Wounds. Poster presentation at the American Association for the Surgery of Trauma meeting, Atlanta, Georgia, September 2005.
8. Soffer D, Klausner J, et al. The usage and availability of blood products in multiple casualty incidents: the experience of a level I trauma center in Israel. Presented at Western Trauma Association Thirty-Sixth Annual Meeting, Big Sky, Montana, 2006.